

1944-1945

عبد كلية التمريض
أ.د / صلاح محمود مهدي

- 10- In sulphuric acid toxicity patient may die immediately from:

 - neurogenic shock
 - asphyxia
 - collapse

11- The danger of criminal abortion is:

 - air embolism
 - vasovagal shock
 - all the above

12- Hyoid bone is broken mostly in cases of:

 - smothering
 - choking
 - strangulation

13- The exit of fire arm injury is smaller than the inlet in the case of:

 - near firing
 - far firing
 - firing in bone

14- Abrasions are common due to:

 - homicide
 - suicide
 - haemophilia

15- Multiple cut wounds are indicative of:

 - ante mortem wound
 - self inflicted wound
 - contused wound

16- Injury by a shot gun from near distance characterized by:

 - common inlet
 - complete dispersion
 - absence of powder marks

17- Less gaping in a wound is correct in:

 - contused wound
 - stab wound
 - cut wound

18- three days after saving a drowned person he may suffer from:

 - acute heart failure
 - broncho pneumonia
 - haemolysis

19- Histamine shock gives rise to the following type of anoxia:

 - anoxic
 - paralytic
 - stagnant

20- One minute after a severe trauma on the testicles, the pulse rate / min. is usually:

 - 45
 - 100
 - 125

HELWAN UNIVERSITY
FACULTY OF NURSING
FORENSIC MEDICINE AND TOXICOLOGY
FINAL TERM EXA

Time allowed : 2 hours
Date : 13/6/2011

طبيب شرعي (Forensic Medicine)

عبد كريمة التبريد
مدرس / مدرس مساعد

A. - Put (✓) or (X): (20 marks)

- 1-Vitamin B12 is one of the coma cocktail ()
- 2-The first line of treatment of the poisoned patient is the antidote ()
- 3-Burning pain & vomiting are common symptoms of corrosives ()
- 4-Face disfigurement is one of the complications of sulfuric acid ()
- 5-It is contraindicated to do gastric lavage in phenol toxicity ()
- 6-There is no physical dependence in habituation ()
- 7-OP toxicity leads to increase level of acetyl choline ()
- 8 -It is impossible to treat drug addiction ()
- 9- The cause of death in OP toxicity is asphyxia ()
- 10-Corrosives are one of the causes of red asphyxia ()
- 11-There is bridging of tissues at the base of cut wound ()
- 12-The edges of contused wound is regular ()
- 13-Shock is one of the complications of wounds ()
- 14-Fire arm wound is characterized by loss of substance ()
- 15-In Fire arm wound from far distance the inlet is larger than the exit ()
- 16-The rope mark in hanging is incomplete ()
- 17-Diatoms (planktons) are sure sign of drowning ()
- 18-Criminal abortion may be complicated with asphyxia ()
- 19-Strangulation is mostly be homicidal ()
- 20-Cut wound is caused by heavy blunt instrument ()

B - MCQ: (20 marks)

1- Naloxone is the best antidote for:

- a. aspirin
- b. opium
- c. corrosives

2- Potash ingestion may produce the following except:

- a. constipation
- b. shock
- c. edema of glottis

3-Arsenic is re- excreted in the:-

- b. Stomach
- c. Duodenum
- d. descending colon

4- Habituation is characterized by except:

- a. physical dependence
- b. tolerance
- c. detrimental effects on the person himself

5- The action of syrup of ipecac begins after:

- a. 5 – 10 min
- b. 15 – 30 min
- c. 30 – 45 min

6- In O.P. poisoning, asphyxia is due to all the following except:

- a. pulmonary edema
- b. spasm of intercostals muscles
- c. bronchospasm

7-Complications of gastric lavage include the following except:



- a. Aspiration pneumonitis
- b. Perforation of the esophagus but not stomach
- c. Asphyxia

8- In drug abuse, the following occur due to non sterile injections except:

- a. hepatitis
- b. lung abscess
- c. appendicitis

9- Management of comatosed patient includes the following except:

- a. CNS stimulants
- b. Glucose
- c. oxygen



1944-1945

عبدك كلية التمريض
أ.د. / هادي محمد هادي

- 10- In sulphuric acid toxicity patient may die immediately from:

 - neurogenic shock
 - asphyxia
 - collapse

11- The danger of criminal abortion is:

 - air embolism
 - vasovagal shock
 - all the above

12- Hyoid bone is broken mostly in cases of:

 - smothering
 - choking
 - strangulation

13- The exit of fire arm injury is smaller than the inlet in the case of:

 - near firing
 - far firing
 - firing in bone

14- Abrasions are common due to:

 - homicide
 - suicide
 - haemophilia

15- Multiple cut wounds are indicative of:

 - ante mortem wound
 - self inflicted wound
 - contused wound

16- Injury by a shot gun from near distance characterized by:

 - common inlet
 - complete dispersion
 - absence of powder marks

17- Less gaping in a wound is correct in:

 - contused wound
 - stab wound
 - cut wound

18- three days after saving a drowned person he may suffer from:

 - acute heart failure
 - broncho pneumonia
 - haemolysis

19- Histamine shock gives rise to the following type of anoxia:

 - anoxic
 - paralytic
 - stagnant

20- One minute after a severe trauma on the testicles, the pulse rate / min. is usually:

 - 45
 - 100
 - 125